

Stanford University Department of Mathematics

Math 53, Fall 2007 — Final Exam

Instructor : Samuel Lisi

Date: 10 December 2007

Duration: 3 hours

FAMILY NAME : _____

GIVEN NAME(S) : _____

STUDENT NUMBER : _____

YOUR SIGNATURE: _____

DO NOT OPEN THIS TEST UNTIL INSTRUCTED TO DO SO.

INSTRUCTIONS:

- Your signature above indicates that you have abided by the Stanford Honor Code while writing this test.
- There are eight questions. The last part of question 8 is a bonus.
- You may quote theorems from your textbook or from class if you make an appropriate reference.
- Show all your work.
- No electronic devices of any kind (e.g. calculators, cell-phones) are allowed.
- There is a table of Laplace transform identities on the last page of the exam. You may use these identities without proof, unless the question indicates otherwise.

Question	Marks
1 [10 pts]	
2 [10 pts]	
3 [15 pts]	
4 [15 pts]	
5 [15 pts]	
6 [15 pts]	
7 [15 pts]	
8 [20 pts]	
Total [100 points]	

1. (a) i. Find a solution to the initial value problem

[10 pts]

$$y' = ty^2, \quad y(0) = 1.$$

- ii. What is the interval of existence of this solution?
iii. Is this solution unique? If yes, explain why. If no, provide a second solution.
- (b) Find the general solution to the differential equation

$$y' + 2ty = e^{-t^2} \sin(t).$$

2. Find the integral curve of

[10 pts]

$$\alpha = (\sin(x + y) + 2x)dx + (y^2 + \sin(x + y))dy$$

that passes through the point $(\pi, 0)$. (You may define this curve implicitly.)

3. (a) Solve the initial value problem :

[15 pts]

$$y'' + 4y = 2 \cos(2t), \quad y(0) = 0, y'(0) = 0.$$

- (b) Solve the initial value problem :

$$y'' + 4y = 2 \cos(2t), \quad y(0) = 1, y'(0) = -1.$$

[15 pts]

4. Let $A = \begin{pmatrix} 2 & 2 & 1 \\ 0 & 2 & 1 \\ 0 & 0 & 2 \end{pmatrix}$.

(a) Find the matrix exponential e^{At} .

(b) Solve the initial value problem :

$$\mathbf{y}' = A\mathbf{y} + \begin{pmatrix} e^{2t} \sin(t) \\ e^{2t} \\ 0 \end{pmatrix}, \quad \mathbf{y}(0) = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}.$$

[15 pts]

5. (a) Sketch the phase portrait for $\mathbf{y}' = \begin{pmatrix} -1 & 2 \\ 2 & -1 \end{pmatrix} \mathbf{y}$.

(b) Determine whether $\mathbf{0}$ is asymptotically stable, stable or unstable.

6. Consider the nonlinear system of differential equations given by :

[15 pts]

$$x' = 2x(2 - x)$$

$$y' = y(1 + x^2)$$

- (a) Find the equilibrium solutions.
- (b) For each equilibrium solution you found in (a), determine its stability.

7. (a) Find the Laplace transform of the solution to the initial value problem [15 pts]

$$y'' + 2y' - y = \cos(2t), \quad y'(0) = 1, y(0) = -1.$$

(You do **not** need to solve for $y(t)$.)

- (b) Find the inverse Laplace transform of $\frac{-4}{(s-1)^2(s-3)}$.

The two parts of this question are unrelated.

8. Consider the nonlinear system of differential equations given by :

[a–c: 10
pts]

$$x' = 2y$$

$$y' = -2x - 4x^3$$

- (a) Show that $(0, 0)$ is the only equilibrium solution.
- (b) What does the linearization at $(0, 0)$ tell us about stability?
- (c) Suppose that $(x(t), y(t))$ is a solution to the nonlinear system. Show that it is an integral curve of

$$\alpha = (2x + 4x^3)dx + 2ydy.$$

- (d) [Bonus – 10 pts] Use the fact that α is exact to conclude that $(0, 0)$ is stable, but not asymptotically stable.

This page has been left blank for your rough work.

This page has been left blank for your rough work.

This page has been left blank for your rough work.

This page has been left blank for your rough work.

Laplace transform identities

In the following, let f be piecewise continuous of exponential order, and $F = \mathcal{L}(f)$:

$$\mathcal{L}(e^{ct} f(t)) = F(s - c)$$

$$\mathcal{L}(tf(t)) = -F'(s)$$

$$\mathcal{L}(1) = \frac{1}{s} \quad \text{for } s > 0$$

$$\mathcal{L}(t^n) = \frac{n!}{s^{n+1}} \quad \text{for } s > 0$$

$$\mathcal{L}(\sin(at)) = \frac{a}{s^2 + a^2} \quad \text{for } s > 0$$

$$\mathcal{L}(\cos(at)) = \frac{s}{s^2 + a^2} \quad \text{for } s > 0$$

$$\mathcal{L}(e^{at}) = \frac{1}{s - a} \quad \text{for } s > a$$

$$\mathcal{L}(e^{at} \sin(bt)) = \frac{b}{(s - a)^2 + b^2} \quad \text{for } s > a$$

$$\mathcal{L}(e^{at} \cos(bt)) = \frac{s - a}{(s - a)^2 + b^2} \quad \text{for } s > a$$

$$\mathcal{L}(t^n e^{at}) = \frac{n!}{(s - a)^{n+1}} \quad \text{for } s > a$$

If f is piecewise differentiable, and both f and f' are of exponential order, then there exists $a > 0$ so that :

$$\mathcal{L}(f'(t)) = sF(s) - f(0) \quad \text{for } s > a.$$